

Characterisation of Dual Nurr1 Activator/DHODH Inhibitor Vidofludimus Calcium and Development Towards a Nurr1 Selective Tool Compound

C. Gege¹, H. Wu², J. Vietor³, T. Wulff¹, M. Jafari¹, E. Peelen¹, T. Stiller³, R. Busch³, J. A. Marschner³, D. Merk³, Z. Sun², H. Kohlhof¹, D. Vitt¹

¹ Immunic AG, Gräfelfing, Germany

² Department of Immunology & Theranostics, Arthur Riggs Diabetes & Metabolism Research Institute, Beckman Research Institute of the City of Hope, Duarte, CA 91010, USA


³ Department of Pharmacy, Ludwig-Maximilians-Universität (LMU) München, 81377 Munich, Germany



Background

Nurr1 Activator

- Protecting neurons from cell death
- Continuous effect independent from focal inflammation



DHODH Inhibitor

- Selective anti-inflammatory effect reduces focal inflammation
- Antiviral effect prevents reactivation of EBV and could stop cross reactive immune responses

Fig. 1: The immunomodulatory and antiviral activity of the Dihydroorotate Dehydrogenase (DHODH) inhibitor vidofludimus calcium (VidoCa) is well known [1]. DHODH catalyzes the rate-limiting step of *de novo* pyrimidine synthesis. We recently found that VidoCa also activates the neuroprotective transcription factor Nuclear Receptor Related 1 (Nurr1), which is an emerging target in neurodegenerative diseases like Parkinson's disease (PD), Alzheimer's disease (AD) or multiple sclerosis (MS) [2].

VidoCa is a potent Nurr1 agonist




Fig. 2: (a) VidoCa activates Gal4-Nurr1 with an EC₅₀ value of 0.4±0.2 μM with a 3.1±0.4-fold maximal activation. Mean±SEM, n ≥3. (b) Binding of VidoCa to the Nurr1 ligand binding domain was confirmed by isothermal titration calorimetry with a K_d value of 0.7 μM. (c) Activity of other DHODH inhibitors related to VidoCa on Nurr1 in the hybrid Gal4-Nurr1 and a full-length Nurr1 (NBRE) reporter gene assay tested at 10 μM. Mean±SEM, n ≥3.

VidoCa is effective in a mouse neuronal cell line




Fig. 3: (a) Nurr1 activation can block neuroinflammation and reactive oxygen species in neuronal cells. (b) VidoCa activates luciferase in a NBRE reporter gene assay activity in the mouse neuronal cell line N2A, with no apparent toxicity at these doses after 48 h treatment. Mean±SEM; * p < 0.05, *** p < 0.001, **** p < 0.0001. (c,d) VidoCa activates Nurr1 target genes like tyrosine hydroxylase (TH) and vesicular amino acid transporter 2 (VMAT2). Mean±SEM; * p < 0.05. (e) VidoCa prevents apoptosis in N2A cells after TNFα/cycloheximide (CHX) stimulation. ##### p < 0.0001 vs. non-treated, **** p < 0.0001 vs. treated DMSO.

VidoCa is effective in human cell lines




Fig. 4: (a) VidoCa remarkably activates Nurr1 target gene VMAT2 in human HMC3 microglial cells. Mean±SEM; ** p < 0.01, *** p < 0.001. (b) VidoCa activates NBRE luciferase activity in the human SH-SY5Y neuronal cell line (with no apparent toxicity at these doses after 48 h treatment). Mean±SEM; ** p < 0.01. (c) VidoCa prevents apoptosis in SH-SY5Y cells after TNFα/cycloheximide (CHX) stimulation. ##### p < 0.0001 vs. non-treated, *** p < 0.001, **** p < 0.0001 vs. treated DMSO.

VidoCa shows Nurr1 related activity in mouse EAE




Fig. 5: In a pilot experimental autoimmune encephalomyelitis model (female C57BL/6 mice; 30 or 150 mg/kg VidoCa p.o./p.d.) mRNA levels were determined: (a) Increased Nurr1 levels were measured in brain tissue upon VidoCa treatment in a dose dependent manner. (b) A significant increase of Nurr1 target genes TH (mRNA, in brain) and (c) brain-derived neurotrophic factor (BDNF) (protein, in heart plasma) were detected upon VidoCa treatment.

VidoCa shows neuroprotective effects in MS patients




Fig. 6: (a) Neurofilament light is used as neurologic biomarker. (b) Improvements in serum NFL were detected in progressive MS patients for VidoCa with a dose of 45 mg (mean change to week 24 compared to placebo in % of baseline).

Optimization towards a Nurr1 selective tool compound




Fig. 7: Compound optimization of deuterated VidoCa results in '2054 with improved selectivity towards Nurr1. (a) Pharmacokinetic properties (e.g. brain exposure) and target engagement of d-VidoCa were further improved during a lead-optimization process, resulting in a large chemical space with new IP. (b) *In vitro* enzyme assays revealed an improved selectivity towards Nurr1 and improved brain penetration for tool compound '2054.

Conclusions and outlook

Conclusion:

- Identified neuroprotective transcription factor Nurr1 as new target for our drug candidate VidoCa
- VidoCa binds to and activates Nurr1
- VidoCa is effective in mouse and human neuronal and microglial cell lines
- VidoCa shows beneficial effects in a murine EAE model, indicating Nurr1 target engagement *in vivo*
- Nurr1 activation might explain the beneficial neuroprotective effects seen in MS patients
- Lead optimization yielded Nurr1 agonist '2054 with improved potency, lack of DHODH inhibitor activity and optimized brain penetration properties

Next steps:

- Additional profiling, e.g. selectivity towards related nuclear receptors
- Use '2054 or related compounds as chemical tools to dissect the DHODH effects from the Nurr1 effects *in vitro* and *in vivo*
- Find improved compounds within the new chemical space
- Elucidate the utility of Nurr1 in other diseases beyond multiple sclerosis

References:

- [1] A. Muehler, et al. Mult. Scler. Relat. Disord. 2020, 43, 102129.
- [2] J. Vietor, et al. J. Med. Chem. 2023, 66, 6391.
- [3] M. Khalil, et al. Nat. Rev. Neurol. 2018, 14, 577.
- [4] A. Abdelhak, et al. JAMA Neurol. 2023, 80, 1317.